

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 July 2004 (29.07.2004)

PCT

(10) International Publication Number
WO 2004/062655 A1

- (51) International Patent Classification⁷: **A61K 31/165**,
A61P 25/06
- (21) International Application Number:
PCT/EP2003/012889
- (22) International Filing Date:
18 November 2003 (18.11.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
03000921.1 16 January 2003 (16.01.2003) EP
- (71) Applicant (for all designated States except US):
NEWRON PHARMACEUTICALS SPA [IT/IT];
Via Ariosto, 21, I-20091 Bresso (IT).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **SALVATI, Patricia**
[IT/IT]; Via Valera, 16, I-20020 Arese (IT). **CALABRESI,**
Marcello [IT/IT]; Via Rogorazza, 29/D, I-21010 Cardano
al Campo (IT). **DHO, Luciano** [IT/IT]; Via de Gasperi,
48, I-20015 Parabiago (IT). **VENERONI, Orietta** [IT/IT];
Via Stephenson, 25, I-20019 Settimo Milanese (IT). **MEL-**
LONI, Piero [IT/IT]; Via Centurelli, 1, I-20091 Bresso
(IT).
- (74) Agents: **MANCINI, Vincenzo** et al.; Ing. A. Giambro-
cono & C. S.r.l., Via Rosolino Pilo, 19/B, I-20129 Milan
(IT).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR,
CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU,
SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ALPHA-AMINOAMIDE DERIVATIVES USEFUL AS ANTIMIGRAINE AGENTS

(57) Abstract: α -Aminoamide derivatives useful as antimigraine agents, particularly for the treatment of head pain conditions such as migraine, cluster headache or other severe headache, are disclosed. The antimigraine agents of the invention are able to reduce or even stop the pain deriving from such conditions without, virtually, any side effects.

WO 2004/062655 A1

ALPHA-AMINOAMIDE DERIVATIVES USEFUL AS ANTIMIGRAINE AGENTS

The invention relates to α -aminoamide derivatives useful as antimigraine agents; particularly, the invention concerns the use of α -aminoamide derivatives in the manufacture of a medicament for the treatment of head pain conditions such as migraine, cluster headache or other severe headache.

The International Headache Classification divides head pain conditions basically into two major categories: primary and secondary headaches.

Primary headache disorders include, for instance, migraine, headache, tension type headache and cluster headache.

Secondary headache disorders result basically from other organic disturbances, such as infection, metabolic disorders, or other systemic illness.

Migraine headaches (also referred to simply as migraines) and cluster headaches are well known medical conditions. Extensive background information on them is contained in references such as "Headache in Clinical Practice" (edited by Silberstein S. et al., Oxford Univ. Press, 1998); and "Headache Disorders: A Management Guide for Practitioners", by Rapoport A. and Sheftell F. (Saunders W.B., Philadelphia, 1996). Various definitions, categories, and diagnostic standards which relate to migraine headaches (and to cluster headaches, described below, and other types of headaches as well) are defined by standardized criteria that were approved and issued by the International Headache Society (IHS), and were published as a supplement to the journal Cephalalgia in 1988.

Despite the emergence of the "triptan" drugs, known to act directly on vasoconstriction, including sumatriptan (also used to treat cluster headaches), naratriptan, zolmitriptan, and rizatriptan, there are no adequately safe, rapid, reliable, and satisfactory treatments for both primary headache disorders, such as recurrent migraines and cluster headaches, and secondary headache disorders such as the ones deriving from infection, metabolic disorders, or other systemic illnesses.

The problems and limitations that plague the treatments contemplating the use of triptans (as well as other known treatments such as, for instance, those comprising the administration of ergotamine analogs etc.) may derive from a number of causes; typically, the main drawbacks include: (i) patients with various types of cardiac or vascular problems cannot take triptan drugs safely; (ii) significant numbers of patients, who repeatedly take any single treatment, run a substantial risk of developing a form of tolerance which can lead to elevated chronic and even continuous headaches; (iii) relief often takes well over half an hour to reach appreciable levels; and (iv) immediately after a treatment, a patient often needs to rest quietly for several hours, which renders it very difficult or impossible for him or her to return to work or get anything else constructive done that day.

WO98/25594 discloses the use of 3,5-diamino-6-(2,3-dichlorophenyl)-1,2,4-triazine, an anticonvulsant also known as lamotrigine, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for use in the treatment and/or prevention of migraine.

WO99/26614 discloses substituted 2-aminoacetamides compounds, preferably α, α' -di-substituted, for treating or ameliorating pain; in view of the activity of said compounds as blockers of sodium channels, the use thereof for the treatment of a number of diseases and conditions mediated by sodium ion influx, among which migraine headache is cited, is therein disclosed too.

It is however known that headache resulting from neurovascular mechanisms does not satisfactorily respond to treatment with analgesic agents and, conversely, antimigraine compounds, such as triptans, do not possess general analgesic properties (Steiner T.J., Findley L.J., Yuen A.W.: "Lamotrigine versus placebo in the prophylaxis of migraine with and without aura", Cephalgia 1997, 17: 109-12; Saxena P.R., Den Boer M.O.: "Pharmacology of antimigraine drugs", J. Neurology 1991, 238 Suppl. 1: S28-35); therefore, the antimigraine properties

cannot be predicted on the basis of pain models.

Further, the Applicant found that most of the substituted 2-aminoacetamides compounds disclosed in WO99/26614 are not effective antimigraine agents and that therefore their usefulness for the treatment of migraine and, generally speaking, of diseases and conditions mediated by sodium ion influx cannot be reasonably predicted in view of their having been recognised as blockers of sodium channels.

Given that antimigraine properties can be predicted neither on the basis of pain models nor in view of the sodium channels blocking activity of compounds, a vast medical need still exists for improved medical treatments which can provide rapid relief from the primary and secondary headaches disorders, particularly from the intense pain of acute migraine or cluster headaches, and which are not associated with problems of migraine recurrence, lingering sedation, unwanted side effects, or elevated health risks such as for patients with cardiac or vascular problems.


The main object of the present invention is to disclose and provide a rapid and highly effective method for treating primary headache disorders including migraine, such as tension type headache, transformed migraine or evolutive headache and cluster headache, as well as secondary headache disorders such as the ones deriving from infection, metabolic disorders, or other systemic illnesses and other acute headaches, in a manner which provides a highly effective treatment with virtually no adverse side effects or lingering after effects (such as drowsiness, grogginess, disorientation, nausea, or the like), thereby allowing the patient to be ready and able to drive, work, or carry out any other normal activity within an hour after that such treatment is commenced.

A further object of the present invention is to disclose and provide a method for treating chronic and/or intractable pain conditions such as, f.i., trigeminal facial pain, chronic paroxysmal hemicrania and the like, resulting from a worsening of the above mentioned primary and secondary headache.

In more general terms, all types of the above mentioned headache disorders, which can be treated according to this invention, are hereinafter collectively referred to as "head pain conditions".

The word "treatment" or the expression "treatment of a condition", whenever employed in this specification, mean inhibiting such condition, i.e. either arresting its development or relieving it or causing its regression as well as preventing its development as soon as the symptoms which are characteristic of such condition begin to appear.

It has now been found that these and other objects of the invention, which will be apparent by the full reading and understanding of the following description, can be attained by the use of an α -aminoamide of formula (I):



wherein:

A is a $-(\text{CH}_2)_m-$ or $-(\text{CH}_2)_n-\text{X}-$, wherein m is 1 or 2; n is zero, 1 or 2; and X is $-\text{O}-$, $-\text{S}-$ or $-\text{NH}-$;

R is a furyl, thienyl, or pyridyl ring or a phenyl ring, unsubstituted or substituted by one or two substituents independently selected from halogen, hydroxy, C_1 - C_4 alkyl, C_1 - C_3 alkoxy and trifluoromethyl;

R_1 is hydrogen or C_1 - C_3 alkyl;

R_2 is hydrogen or C_1 - C_2 alkyl, unsubstituted or substituted by hydroxy or phenyl; phenyl, unsubstituted or substituted by one or two substituents independently selected from C_1 - C_3 alkyl, halogen, hydroxy, C_1 - C_2 alkoxy or trifluoromethyl;

R_3 is hydrogen or C_1 - C_3 alkyl;

if the case, either as a single isomer, or as a mixture thereof, or a pharmaceutically acceptable derivative thereof;

in the manufacture of a medicament for the treatment of head pain conditions.

The alkyl and alkoxy groups may be branched or straight

chain groups. A halogen atom is preferably fluorine, chlorine or bromine, in particular fluorine or chlorine.

A C₁-C₄ alkyl is a linear or branched alkyl group; preferably a methyl, ethyl, propyl, isopropyl, butyl or tert-butyl; most preferably, it is a methyl or ethyl group.

A C₁-C₃ alkoxy is a linear or branched alkoxy group; preferably a methoxy, ethoxy, propoxy, isopropoxy; most preferably, it is a methoxy or ethoxy group.

A thienyl ring is, for instance, a 2- or 3-thienyl ring.

A pyridyl ring is, for instance, a 2-, or 3- or 4-pyridyl, in particular a 3-pyridyl ring.

A furyl ring is, for instance, a 2- or 3-furyl ring.

A substituted phenyl ring is preferably substituted by one or two substituents chosen independently from halogen, C₁-C₃ alkyl and trifluoromethyl.

A C₁-C₂ alkyl group substituted by hydroxy is preferably a hydroxymethyl or 1-hydroxyethyl group.

A C₁-C₂ alkyl group substituted by a phenyl ring is preferably a benzyl or phenethyl group.

Preferably, in formula (I) above:

A is a group selected from -CH₂-CH₂-, -CH₂-O-, -CH₂-S-, -CH₂-CH₂-O-;

R is a phenyl ring, unsubstituted or substituted by one or two substituents independently selected from halogen, C₁-C₃ alkyl or a methoxy group; or a thienyl ring;

R₁ is hydrogen or C₁-C₂ alkyl;

R₂ is hydrogen or methyl, unsubstituted or substituted by hydroxy, or phenyl unsubstituted or substituted by C₁-C₂ alkyl, halogen, hydroxy, methoxy or trifluoromethyl; and

R₃ is hydrogen or C₁-C₂ alkyl.

A further preferred group of α -aminoamides comprises the compounds of formula (I) above wherein:

A is -CH₂-O-, -CH₂-S- or -CH₂-CH₂-;

R is a phenyl ring, unsubstituted or substituted by one or two halogen atoms;

R₁ is hydrogen;

R₂ is hydrogen or methyl, unsubstituted or substituted by hydroxy or phenyl ring, unsubstituted or substituted by a halogen atom;

R₃ is hydrogen or methyl.

Preferred α -aminoamides for the present invention are:

- 2-(4-benzyloxybenzylamino)propanamide;
- 2-[4-(2-fluorobenzyloxy)benzylamino]propanamide;
- 2-[4-(2-chlorobenzyloxy)benzylamino]propanamide;
- 2-[4-(3-fluorobenzyloxy)benzylamino]propanamide;
- 2-[4-(3-chlorobenzyloxy)benzylamino]propanamide;
- 2-[4-(4-fluorobenzyloxy)benzylamino]propanamide;
- 2-[4-(2-fluorobenzyloxy)benzylamino]-N-methyl-propanamide;
- 2-[4-(3-fluorobenzyloxy)benzylamino]-N-methyl-propanamide;
- 2-[4-(2-fluorobenzyloxy)benzylamino]-3-hydroxy-propanamide;
- 2-[4-(3-fluorobenzyloxy)benzylamino]-3-hydroxy-propanamide;
- 2-(4-benzyloxybenzylamino)-3-hydroxy-N-methylpropanamide;
- 2-[4-(2-fluorobenzyloxy)benzylamino]-3-hydroxy-N-methylpropanamide;
- 2-[4-(2-chlorobenzyloxy)benzylamino]-3-hydroxy-N-methylpropanamide;
- 2-[4-(3-fluorobenzyloxy)benzylamino]-3-hydroxy-N-methylpropanamide;
- 2-[4-(3-chlorobenzyloxy)benzylamino]-3-hydroxy-N-methylpropanamide;
- 2-(4-(2-thienylmethylenoxy)benzylamino)-propanamide;
- 2-[4-(2-(3-fluorophenyl)ethyl)benzylamino]-propanamide;
- 2-[4-benzylthiobenzylamino]-propanamide;
- 2-[4-benzyloxybenzylamino]-3-phenyl-N-methylpropanamide;
- 2-[4-benzyloxybenzylamino]-N-methylbutanamide;
- 2-[4-benzyloxybenzylamino]-2-phenyl-acetamide;
- 2-[4-(2-fluorobenzyloxy)benzylamino]-2-phenyl-acetamide;
- 2-[4-(3-fluorobenzyloxy)benzylamino]-2-phenyl-acetamide;
- 2-[4-(3-chlorobenzyloxy)benzylamino]-2-phenyl-acetamide;
- 2-[4-(3-fluorobenzyloxy)benzylamino]-2-(2-fluorophenyl)-acetamide;
- 2-[4-(3-fluorobenzyloxy)benzylamino]-2-(3-fluorophenyl)-

acetamide;

2-[4-(3-chlorobenzyloxy)benzylamino]-2-(3-fluorophenyl)-
acetamide;

if the case, either as a single isomer or as a mixture thereof, or a pharmaceutically acceptable derivative thereof.

In the present invention, the following α -aminoamides are most preferred: (S)-(+)-2-[4-(3-fluorobenzyloxy)benzylamino]-propanamide (internal code and hereinafter NW-1015), (S)-(+)-2-[4-(2-fluorobenzyloxy)benzylamino]-propanamide (internal code and hereinafter NW-1029) and (S)-(+)-2-[4-(3-chlorobenzyloxy)benzylamino]-propanamide (internal code and hereinafter NW-1039).

The α -aminoamides of formula (I) and the analgesic activity thereof, in particular against chronic and neuropathic pain in mammals including humans, are disclosed in WO90/14334, WO94/22808, WO97/05102, WO99/26614, WO99/35123 and WO99/35125; any of the α -aminoamides of the above formula (I) can be prepared according to what disclosed in said documents which are herein incorporated by reference as far as the preparation of said α -aminoamides is concerned.

WO90/14334, WO94/22808, WO97/05102, WO97/05111, disclose substituted benzylaminoamide compounds active on the central nervous system and useful as anti-epileptic, anti-Parkinson, neuroprotective, antidepressant, antispastic hypnotic agents (Pevarello P., Bonsignori A., Doster P., Heidempergher F., Pincioli V., Colombo M., McArthur R.A., Salvati P., Post C., Fariello R.G. and Varasi M.: "Synthesis and anticonvulsant activity of a new class of 2-[(arylalkyl)amino]alkanamide derivatives", J. Med. Chemistry 1998, 41: 579-590).

WO99/35123 and WO99/35125 disclose substituted benzylaminopropanamide compounds active on the central nervous system and useful as analgesic agents. It has been shown that the α -aminoamides disclosed in WO99/35123 and WO99/35125 are Na⁺ channel blockers having analgesic effects against some models of pain in animals (Faravelli L., Maj R., Veneroni O., Fariello R.G., Benatti L., Salvati P., Society for Neuroscience

2000; 26 (1): 1218).

A further aspect of the invention relates to a method for the treatment of head pain conditions in a mammal, including humans, in need thereof comprising administering to the mammal a therapeutically effective dose of at least one α -aminoamide of formula (I) as above defined or a pharmaceutically acceptable salt thereof.

Particularly, the mammal in need of the above mentioned treatment is administered a dose of the α -aminoamide of formula (I) as above defined which ranges from about 0.05 to 20 mg/kg body weight per day; preferably in the range of about 0.5 to 10 mg/kg day; most preferably in the range of about 0.5 to 5 mg/kg day.

Head pain conditions in a mammal, including humans, can thus be inhibited, alleviated and prevented. Examples of head pain conditions in mammals which can be treated by the above defined α -aminoamide of formula (I) are those involving a cerebral vasodilatation mechanism and include both primary and secondary headache disorders; particularly, those primary headache disorders which derive from the intense pain of acute migraine or cluster headaches or from vascular mechanisms and those secondary headache disorders which derive from infection, metabolic disorders, or other systemic illnesses.

In particular, examples of head pain conditions that can be treated by the α -aminoamide of formula (I) as above defined include migraine such as, for instance, acute, transformed or vascular migraine; headache such as, for instance, acute, cluster, evolutive or tension type headache; neuralgia such as, for instance, trigeminal neuralgia; hemicrania such as, for instance, chronic paroxysmal hemicrania; facial pain and arachnoiditis.

Further, a "pharmaceutically acceptable derivative" of the α -aminoamide of formula (I) as above defined is herein meant to include any pharmaceutically acceptable metabolite, bioprecursor and/or pro-drug, i.e. a compound which has a

structural formula different from the one of the α -aminoamide of formula (I) as above defined and yet is directly or indirectly converted *in vivo* into a compound having their structural formula, upon administration to a mammal, particularly a human being.

Examples of pharmaceutically acceptable derivatives of the α -aminoamide of formula (I) as above defined include acid addition salts with inorganic acids, e.g. nitric, hydrochloric, hydrobromic, sulphuric and phosphoric acids and the like, or organic acids, e.g. acetic, propionic, glycolic, lactic, malonic, malic, tartaric, citric, succinic, benzoic, cinnamic, mandelic, methanesulfonic, p-toluenesulfonic and salicylic acids and the like.

The α -aminoamide of formula (I) as above defined can be administered as the "active ingredient" of a pharmaceutically acceptable composition which can be prepared by conventional procedures known in the art, for instance by mixing the active ingredient with pharmaceutically acceptable, therapeutically inert organic and/or inorganic carrier materials.

The composition comprising the above defined α -aminoamide can be administered in a variety of dosage forms, e.g. orally, in the form of tablets, troches, capsules, sugar or film coated tablets, liquid solutions, emulsions or suspensions; rectally, in the form of suppositories; parenterally, e.g. by intramuscular or intravenous injection or infusion; and transdermally.

Suitable pharmaceutically acceptable, therapeutically inert organic and/or inorganic carrier materials useful in the preparation of such composition include, for example, water, gelatin, gum arabic, lactose, starch, cellulose, magnesium stearate, talc, vegetable oils polyalkyleneglycols and the like. The composition comprising the α -aminoamide of formula (I) as above defined can be sterilized and may contain further components, well known to the skilled in the art, such as, for example, preservatives, stabilizers, wetting or emulsifying agents, e.g. paraffin oil, mannide monooleate, salts to adjust

osmotic pressure, buffers and the like.

For example, the solid oral forms may contain, together with the active ingredient, diluents, e.g. lactose, dextrose, saccharose, cellulose, corn starch or potato starch; lubricants, e.g. silica, talc, stearic acid, magnesium or calcium stearate, and/or polyethylene glycols; binding agents, e.g. starches, arabic gums, gelatin, methylcellulose, carboxymethylcellulose or polyvinyl pyrrolidone; disgregating agents, e.g. a starch, alginic acid, alginates or sodium starch glycolate; effervescing mixtures; dyestuffs; sweeteners; wetting agents such as lecithin, polysorbates, laurylsulphates; and, in general, non-toxic and pharmacologically inactive substances used in pharmaceutical formulations. Said pharmaceutical preparations may be manufactured in known manner, for example, by means of mixing, granulating, tableting, sugar-coating, or film-coating processes.

The oral formulations comprise sustained release formulations that can be prepared in conventional manner, for instance by applying an enteric coating to tablets and granules.

The liquid dispersion for oral administration may be e.g. syrups, emulsions and suspension.

The syrups may contain as carrier, for example, saccharose or saccharose with glycerine and/or mannitol and/or sorbitol.

Suspensions and emulsions may contain as a carrier, for example, a natural gum, agar, sodium alginate, pectin, methylcellulose, carboxymethylcellulose, or polyvinyl alcohol. The suspensions or solutions for intramuscular injections may contain, together with the active compound, a pharmaceutically acceptable carrier, e.g. sterile water, olive oil, ethyl oleate, glycols, e.g. propylene glycol, and, if desired, a suitable amount of lidocaine hydrochloride. The solutions for intravenous injections or infusion may contain as carrier, for example, sterile water or preferably they may be in the form of sterile, aqueous, isotonic saline solutions.

The suppositories may contain, together with the active

ingredient, a pharmaceutically acceptable carrier, e.g. cocoa butter, polyethylene glycol, a polyoxyethylene sorbitan fatty acid ester surfactant or lecithin.

The composition comprising the α -aminoamide of formula (I) as above defined is generally in the form of a dose unit containing, for example, 35 to 350 mg of active ingredient per unit dosage form.

Suitable treatment is given 1, 2 or 3 times daily, depending upon clearance rate. Accordingly, the desired dose may be presented in a single dose or as divided doses administered at appropriate intervals, for example two to four or more sub-doses per day. Treatment is preferably commenced before the onset of a migraine episode and continued indefinitely.

The pharmaceutical compositions comprising the α -aminoamide of formula (I) as above defined will contain, per dosage unit, e.g., capsule, tablet, powder injection, teaspoonful, suppository and the like from about 35 to about 350 mg of the active ingredient.

Optimal therapeutically effective doses to be administered may be readily determined by those skilled in the art and will vary, basically, with the strength of the preparation, with the mode of administration and with the advancement of the condition or disorder treated. In addition, factors associated with the particular subject being treated, including subject age, weight, diet and time of administration, will result in the need to adjust the dose to an appropriate therapeutically effective level.

The advantages deriving from the use and the method of the invention as above defined are many, and include the possibility to prevent and treat basically all types of migraine, cluster or other severe headaches and to reduce or entirely stop pain caused by the headaches in a rapid and highly effective way; specifically both primary and secondary headache disorders or other systemic illnesses and other acute headaches.

Besides, the use and method of the invention did not show, virtually, any adverse side effects or lingering after effects (such as drowsiness, grogginess, disorientation, nausea, or other such problems), thereby allowing the patient to be ready and able to drive, work, or carry out any other normal activity within an hour after that such treatment is commenced.

The following examples illustrate the invention without limiting it.

EXAMPLE 1

The following pharmaceutical composition, as well as the ones of Examples 2 and 3, comprising an α -aminoamide of formula (I) as above defined, were prepared by mixing the ingredients below listed, employing methods usual in the pharmaceutical field and known to the skilled in the art.

One 35-mg capsule contains

NW-1015	35.00 mg
Crospovidone	4.37 mg
Microcrystalline cellulose	5.95 mg
Magnesium stearate	0.17 mg
Colloidal silicon dioxide	0.18 mg

EXAMPLE 2

One 100-mg capsule contains

NW-1015	100.00 mg
Crospovidone	7.50 mg
Microcrystalline cellulose	8.95 mg
Magnesium stearate	1.50 mg
Colloidal silicon dioxide	0.30 mg

EXAMPLE 3

One 175-mg capsule contains

NW-1015	175.00 mg
Crospovidone	13.05 mg
Microcrystalline cellulose	15.57 mg
Magnesium stearate	2.61 mg
Colloidal silicon dioxide	0.49 mg

PHARMACOLOGY

The antimigraine activity of the above defined α -aminoamide

of formula (I) has been proven by the studies reported hereinafter of vascular migraines and related disorders on animal models.

Studies [Reuter U., Sanchez del Rio M., Moskowitz M.A., "Experimental models of migraine", Functional neurology (15) Suppl. 3 9-18, 2000; Magnus L., "Nonepileptic uses of gabapentin. Experience", Epilepsia, 1999; 40 (Suppl. 6) S66-72; discussion S73-S74; Peitl B., Pethô G., Pórszász R., Németh J. and Szolcsányi J., "Capsaicin-insensitive sensory-efferent meningeal vasodilatation evoked by electrical stimulation of trigeminal nerve fibres in the rat"] of the regional cortical blood flow (CBF) have shown that during the headache phase of migraine, a dilatation of both large extra and intracranial arteries occurs. This event has been associated to an antidromic activation with a consequent vasoactive neuropeptides release, and to an orthodromic activation of the trigeminal fibers associated with an increased neuronal activity.

The above defined α -aminoamides have been found to be active in inhibiting cerebral vasodilatation evoked by electrical stimulation of ophthalmic branch of rat trigeminal ganglion and are therefore deemed to be useful as antimigraine agents.

GENERAL METHODS

Animals and surgery

Male Wistar rats (250-350 g) were anesthetized with sodium pentobarbital (50 mg/kg i.p.) dissolved in saline.

The trachea and left femoral artery were cannulated for artificial ventilation (55 strokes/min) and for measurement of mean blood pressure (MBP) respectively. The femoral vein was cannulated for the intravenous administration of test agents. Body temperature was maintained at 37-38°C by automatic control of a heating pad.

Animals were placed in a stereotaxic frame and a longitudinal incision was made in the scalp. A burr hole was drilled in the skull and a stainless steel bipolar electrode

(Plastic One MS 306) was lowered into left ophthalmic branch of the trigeminal ganglion (3.8 mm dorsal to bregma, 2.5 mm lateral from the midline and 9.5 mm below the dural surface) and secured with dental cement.

Correct placement of the electrode was confirmed by a brief electrical stimulation, which cause movement of the jaw due to activation of the trigeminal fiber. Following removal of the brain, the correct position of the electrode into the fiber, was visually checked at the end of each experiment.

A second hole was drilled ipsilateral of the electrode (1.5 mm rostral to bregma, and 1.5 mm lateral from the sagittal suture) and a needle probe (tip diameter 0.8 mm) of a laser doppler flowmeter was fixed pointing with its tip onto a branch of the middle cerebral artery (MCA) and Cerebral Blood Flow (CBF) change recorded on-line by the PeriFlux 4001 Laser Doppler system.

Artefacts of the laser Doppler reading during electrical stimulation of the trigeminal ganglion due to muscular movements were prevented by a bolus of i.v. injection of the neuromuscular blocker pancuronium bromide (0.6 mg/kg i.v.). Anesthesia and neuromuscular blockade were maintained all over the experiment with an infusion of sodium pentobarbital and pancuronium (12.5 mg/kg/h + 2.4 mg/kg/h, respectively).

EXPERIMENTAL PROTOCOL

At the end of the surgery, a pause of thirty minutes was taken in order to stabilize the measured parameters.

Rest CBF was increased by electrical stimulation with rectangular pulse of 0.5 msec length, 1-10 Hz, 0.5-1 mA for periods of 30 seconds. After two averaged pre-drug stimulations, vehicle or drugs were administered.

The α -aminoamide compounds NW-1015, NW-1029, NW-1039, representative of the invention, and the comparative 2-(4-(2-fluorobenzyloxy)benzylamino)-2-methyl-propanamide (internal code and hereinafter NW1050), 2-(4-(4-fluorobenzyloxy)benzylamino)-2-methyl-propanamide (internal code and hereinafter NW1055, also one of the compounds tested

in WO99/26614), 4-(4'-fluorophenoxy)benzaldehyde semicarbazone (hereinafter Co102862, tested in WO00/61188) and vehicle, were administered intravenously after second basal stimulation registering the responses to evoked flow by electrical stimulation at 5, 15, 30 and 60 min after treatment.

NW-1015, was administered at the doses of 1, 2 and 5 mg/kg; NW-1029 was administered at the doses of 5 and 10 mg/kg and NW-1039 at the doses of 2 and 5 mg/kg whereas NW-1050, NW-1055 and Co102862 were all administered at the dose of 20 mg/kg.

The responses obtained after drug administration were compared to the vehicle group, and were reported as a percentage of the inhibition of the evoked CBF. Data were compared by analysis of covariance (ANCOVA) followed by Dunnet's test.

RESULTS

The antimigraine effect of the tested compounds was observed and measured as the percentage of the inhibition after i.v. administration of the above mentioned representative compounds of the CBF evoked in control condition. The data reported in table 1 below show the inhibitory activity of the tested compounds on the CBF response evoked by electrical stimulation of the left ophthalmic branch of the trigeminal ganglion.

Table 1

Compound	Dose (mg/kg)	Time (min) after administration			
		5	15	30	60
% Inhibition of the CBF response					
NW-1015	1	25.6±2.9	25.8±2.7	36.6±4.7**	15.7±9.8
NW-1015	2	15.0±1.8*	22.0±3.8**	20.6±6.2*	4.0±6.3
NW-1015	5	25.2±2.6	35.5±6.5**	34.7±1.9**	27.1±3.6
NW-1029	5	15.5±3.7	20.8±3.5	27.8±3.9**	30.6±2.8
NW-1029	10	23.9±3.9	34.9±8.2**	35.3±2.6**	35.7±3.9**
NW-1039	2	24.2±6.2	31.3±6.2*	32.1±8.1**	36.2±9.1**
NW-1039	5	45.0±6.3**	41.7±1.8**	38.1±2.6**	20.9±9.4
NW-1050	20	1.2±0.3	1.8±0.6	1.7±0.8	0.9±0.1
NW-1055	20	6.9±4.2	3.4±3.2	11.3±2.0	14.0±4.0
Co102862	20	0.9±0.1	1.1±0.3	1.2±0.2	0.7±0.1

* $p < 0.05$, ** $p < 0.01$ by Dunnett's t test

Results (mean \pm S.E. of 4/6 rats per group) are expressed as percent of the inhibition on evoked CBF response versus the vehicle.

NW-1015 showed a long lasting i.v. activity reducing the evoked CBF up to 60 min after administration. NW-1029 equally, reduced the evoked CBF compared to the vehicle up to 60 min post administration. NW-1039 at the doses tested produced a decrease of the evoked CBF response compared to the vehicle group.


NW-1055 at 20 mg/kg caused a light inhibition of the evoked CBF, which however did not reach a statistically significant difference in comparison with the vehicle treated group. NW-1050 and Col02862 at 20 mg/kg were completely inactive.

The above data confirm that representative substituted 2-aminoacetamides compounds disclosed in WO99/26614 are not effective as antimigraine agents and that therefore such activity cannot be reasonably predicted in view of having recognised such compounds as blockers of sodium channels. Particularly, the preferred α, α' -di-substituted acetamides (2-methylpropanamide derivatives) showed a far lower potency than the above defined amides of formula (I).

The above biological activities confirm that the α -aminoamide of formula (I) as above defined can be used as antimigraine agents, in particular, to treat head pain conditions which involve a cerebral vasodilatation mechanism.

CLAIMS

1. Use of an α -aminoamide of formula (I):



wherein:

A is a $-(\text{CH}_2)_m-$ or $-(\text{CH}_2)_n-\text{X}-$, wherein m is 1 or 2; n is zero, 1 or 2; and X is $-\text{O}-$, $-\text{S}-$ or $-\text{NH}-$;

R is a furyl, thienyl, or pyridyl ring or a phenyl ring, unsubstituted or substituted by one or two substituents independently selected from halogen, hydroxy, C_1 - C_4 alkyl, C_1 - C_3 alkoxy and trifluoromethyl;

R_1 is hydrogen or C_1 - C_3 alkyl;

R_2 is hydrogen or C_1 - C_2 alkyl, unsubstituted or substituted by hydroxy or phenyl; phenyl, unsubstituted or substituted by one or two substituents independently selected from C_1 - C_3 alkyl, halogen, hydroxy, C_1 - C_2 alkoxy or trifluoromethyl;

R_3 is hydrogen or C_1 - C_3 alkyl;

if the case, either as a single isomer, or as a mixture thereof, or a pharmaceutically acceptable derivative thereof;

in the manufacture of a medicament for the treatment of head pain conditions.

2. Use of an α -aminoamide according to claim 1, wherein in formula (I):

A is a group selected from $-\text{CH}_2-\text{CH}_2-$, $-\text{CH}_2-\text{O}-$, $-\text{CH}_2-\text{S}-$, $-\text{CH}_2-\text{CH}_2-\text{O}-$;

R is a phenyl ring, unsubstituted or substituted by one or two substituents independently selected from halogen, C_1 - C_3 alkyl or a methoxy group; or a thienyl ring;

R_1 is hydrogen or C_1 - C_2 alkyl;

R_2 is hydrogen or methyl, unsubstituted or substituted by hydroxy, or phenyl unsubstituted or substituted by C_1 - C_2 alkyl, halogen, hydroxy, methoxy or trifluoromethyl; and

R_3 is hydrogen or C_1 - C_2 alkyl.

3. Use of an α -aminoamide according to claim 1 or 2,

wherein in formula (I):

A is $-\text{CH}_2-\text{O}-$, $-\text{CH}_2-\text{S}-$ or $-\text{CH}_2-\text{CH}_2-$;

R is a phenyl ring, unsubstituted or substituted by one or two halogen atoms;

R_1 is hydrogen;

R_2 is hydrogen or methyl, unsubstituted or substituted by hydroxy or phenyl ring, unsubstituted or substituted by a halogen atom; and

R_3 is hydrogen or methyl.

4. Use of an α -aminoamide according to claim 1, wherein the α -aminoamide is selected from:

- 2-(4-benzyloxybenzylamino)propanamide;
- 2-[4-(2-fluorobenzyloxy)benzylamino]propanamide;
- 2-[4-(2-chlorobenzyloxy)benzylamino]propanamide;
- 2-[4-(3-fluorobenzyloxy)benzylamino]propanamide;
- 2-[4-(3-chlorobenzyloxy)benzylamino]propanamide;
- 2-[4-(4-fluorobenzyloxy)benzylamino]propanamide;
- 2-[4-(2-fluorobenzyloxy)benzylamino]-N-methyl-propanamide;
- 2-[4-(3-fluorobenzyloxy)benzylamino]-N-methyl-propanamide;
- 2-[4-(2-fluorobenzyloxy)benzylamino]-3-hydroxy-propanamide;
- 2-[4-(3-fluorobenzyloxy)benzylamino]-3-hydroxy-propanamide;
- 2-(4-benzyloxybenzylamino)-3-hydroxy-N-methylpropanamide;
- 2-[4-(2-fluorobenzyloxy)benzylamino]-3-hydroxy-N-methylpropanamide;
- 2-[4-(2-chlorobenzyloxy)benzylamino]-3-hydroxy-N-methylpropanamide;
- 2-[4-(3-fluorobenzyloxy)benzylamino]-3-hydroxy-N-methylpropanamide;
- 2-[4-(3-chlorobenzyloxy)benzylamino]-3-hydroxy-N-methylpropanamide;
- 2-(4-(2-thienylmethylenoxy)benzylamino)-propanamide;
- 2-[4-(2-(3-fluorophenyl)ethyl)benzylamino]-propanamide;
- 2-[4-benzylthiobenzylamino]-propanamide;
- 2-[4-benzyloxybenzylamino]-3-phenyl-N-methylpropanamide;
- 2-[4-benzyloxybenzylamino] N-methylbutanamide;
- 2-[4-benzyloxybenzylamino]-2-phenyl-acetamide;

2-[4-(2-fluorobenzyloxy)benzylamino]-2-phenyl-acetamide;
2-[4-(3-fluorobenzyloxy)benzylamino]-2-phenyl-acetamide;
2-[4-(3-chlorobenzyloxy)benzylamino]-2-phenyl-acetamide;
2-[4-(3-fluorobenzyloxy)benzylamino]-2-(2-fluorophenyl)-
acetamide;

2-[4-(3-fluorobenzyloxy)benzylamino]-2-(3-fluorophenyl)-
acetamide;

2-[4-(3-chlorobenzyloxy)benzylamino]-2-(3-fluorophenyl)-
acetamide;

if the case, either as a single isomer or as a mixture thereof, or a pharmaceutically acceptable derivative thereof.

5. Use of an α -aminoamide according to any of the previous claims, wherein the α -aminoamide is selected from: (S)-(+)-2-[4-(3-fluorobenzyloxy)benzylamino]-propanamide, (S)-(+)-2-[4-(2-fluorobenzyloxy)benzylamino]-propanamide and (S)-(+)-2-[4-(3-chlorobenzyloxy)benzylamino]-propanamide.

6. Use according to any of the previous claims, wherein the head pain conditions are involving a cerebral vasodilatation mechanism.

7. Use according to any of the previous claims, wherein head pain conditions are both primary and secondary headache disorders.

8. Use according to any of the previous claims, wherein the primary headache disorders derive from the intense pain of acute migraine or cluster headaches or from vascular mechanisms; and the secondary headache disorders derive from infection, metabolic disorders, or other systemic illnesses.

9. Use according to any of the previous claims, wherein head pain conditions include migraine, headache, neuralgia, hemicrania, facial pain and arachnoiditis.

10. Use according to any of the previous claims, wherein migraine is acute, transformed or vascular migraine; headache is acute, cluster, evolutive or tension type headache; neuralgia is trigeminal neuralgia; hemicrania is chronic paroxysmal hemicrania.

11. A method for the treatment of head pain conditions in a

mammal in need thereof comprising administering to the mammal a therapeutically effective dose of at least one α -aminoamide of formula (I) as defined in any of claims 1 to 5.

12. A method according to the previous claim, wherein the mammal is administered a dose of the α -aminoamide of formula (I) as defined in any of claims 1 to 5 which ranges from about 0.05 to 20 mg/kg body weight per day.

13. A method according to claim 11 or 12, wherein the mammal is administered a dose of the α -aminoamide of formula (I) as defined in any of claims 1 to 5 which ranges from about 0.5 to 10 mg/kg day.

14. A method according to any of claims from 11 to 13, wherein the mammal is administered a dose of the α -aminoamide of formula (I) as defined in any of claims 1 to 5 which ranges from about 0.5 to 5 mg/kg day.

15. A method according to any of claims from 11 to 14, wherein the head pain conditions are as defined in any of claims 6 to 10.

INTERNATIONAL SEARCH REPORT

International Application No.

EP 03/12889

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K31/165 A61P25/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 99 35125 A (PEVARELLO PAOLO ;VARASI MARIO (IT); NEWRON PHARM SPA (IT); SALVATI) 15 July 1999 (1999-07-15) cited in the application * page 2, formula I * page 3, line 5 page 12, line 23 - line 26 page 6, line 14 - line 17 ---	1-15
Y	WO 99 26614 A (CAI SUI XIONG ;LAN NANCY C (US); COCENSYS INC (US); WANG YAN (US)) 3 June 1999 (1999-06-03) cited in the application page 8, line 3 page 24, line 12 - line 22 * page 33, formula II * ---	1-15

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

3 February 2004

Date of mailing of the international search report

13/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Albayrak, T

INTERNATIONAL SEARCH REPORT

International Application No

EP 03/12889

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>EP 1 123 702 A (EISAI CO LTD) 16 August 2001 (2001-08-16) page 3, line 9 - line 17 page 14; claim 4</p>	1-15

BEST AVAILABLE COPY

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 03/12889

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9935125	A	15-07-1999	AT 238273 T	15-05-2003
			BR 9814548 A	10-10-2000
			CA 2316902 A1	15-07-1999
			DE 69813896 D1	28-05-2003
			DE 69813896 T2	06-11-2003
			DK 1045830 T3	04-08-2003
			EA 2763 B1	29-08-2002
			WO 9935125 A1	15-07-1999
			EP 1045830 A1	25-10-2000
			ES 2194392 T3	16-11-2003
			HU 0100870 A2	30-07-2001
			JP 2002508302 T	19-03-2002
			NO 20003399 A	02-08-2000
			NZ 505440 A	01-02-2002
			US 6306903 B1	23-10-2001
WO 9926614	A	03-06-1999	AU 749214 B2	20-06-2002
			AU 1421599 A	15-06-1999
			CA 2310664 A1	03-06-1999
			EP 1032377 A1	06-09-2000
			JP 2001523710 T	27-11-2001
			WO 9926614 A1	03-06-1999
			US 2003130295 A1	10-07-2003
			US 6479484 B1	12-11-2002
			US 2002016464 A1	07-02-2002
EP 1123702	A	16-08-2001	CA 2345406 A1	27-04-2000
			EP 1123702 A1	16-08-2001
			CN 1324239 T	28-11-2001
			WO 0023075 A1	27-04-2000
			JP 2000191533 A	11-07-2000

BEST AVAILABLE COPY

BEST AVAILABLE COPY